

THE EXISTENCE OF QUASIMEROMORPHIC MAPPINGS

EMIL SAUCAN

ABSTRACT. We prove that a Kleinian group G acting upon \mathbb{H}^n admits a non-constant G -automorphic function, even if it has torsion elements, provided that the orders of the elliptic (torsion) elements are uniformly bounded. This is accomplished by developing a technique for mashing distinct fat triangulations while preserving fatness.

1. INTRODUCTION

The object of this article is the study of the existence of G -automorphic quasimeromorphic mappings $f : \mathbb{H}^n \rightarrow \widehat{\mathbb{R}^n}$, $\widehat{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$; i.e. such that

$$(1.1) \quad f(g(x)) = f(x) \quad ; \quad \forall x \in \mathbb{H}^n; \forall g \in G ;$$

where G is Kleinian group acting upon \mathbb{H}^n , and where quasimeromorphic mappings are defined as follows:

Definition 1.1. Let $D \subseteq \mathbb{R}^n$ be a domain; $n \geq 2$, and let $f : D \rightarrow \mathbb{R}^m$. f is called *ACL* (*absolutely continuous on lines*) iff:

- (i) f is continuous
- (ii) for any n -interval $Q = \overline{Q} = \{a_i \leq x_i \leq b_i \mid i = 1, \dots, n\}$, f is absolutely continuous on almost every line segment in Q , parallel to the coordinate axes.

Lemma 1.2 ([V], 26.4). *If $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ is ACL, then f admits partial derivatives almost everywhere.*

The result above justifies the following Definition:

Definition 1.3. $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ is *ACL^p* iff its derivatives are locally L^p integrable, $p \geq 1$.

Definition 1.4. Let $D \subseteq \mathbb{R}^n$ be a domain; $n \geq 2$ and let $f : D \rightarrow \mathbb{R}^m$ be a continuous mapping. f is called

- (1) *quasiregular* iff (i) f is *ACLⁿ* and
(ii) $\exists K \geq 1$ s.t.

$$(1.2) \quad |f'(x)| \leq K J_f(x) \text{ a.e.}$$

where $f'(x)$ denotes the formal derivative of f at x , $|f'(x)| = \sup_{|h|=1} |f'(x)h|$,

and where $J_f(x) = \det f'(x)$.

The smallest K that satisfies (4.1) is called the *outer dilatation* of f .

Date: 27.4.2003.

1991 *Mathematics Subject Classification.* 30C65, 57R05, 57M60.

Key words and phrases. quasimeromorphic mapping, fat triangulation.

This paper represents part of the authors Ph.D. Thesis written under the supervision of Prof. Uri Srebro.

- (2) *quasiconformal* iff $f : D \rightarrow f(D)$ is a local homeomorphism.
- (3) *quasimeromorphic* iff $f : D \rightarrow \widehat{\mathbb{R}^n}$, $\widehat{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$ is quasiregular, where the condition of quasiregularity at $f^{-1}(\infty)$ can be checked by conjugation with auxiliary Möbius transformations.

Remark 1.5. One can extend the definitions above to oriented \mathcal{C}^∞ Riemannian n -manifolds by using coordinate charts.

Our principal goal is to prove:

Theorem 1.6. *Let G be a Kleinian group with torsion acting upon \mathbb{H}^n , $n \geq 3$. If the elliptic elements (i.e. torsion elements) of G have uniformly bounded orders, then there exists a non constant G -automorphic quasimeromorphic mapping $f : \mathbb{H}^n \rightarrow \mathbb{R}^n$.*

The question whether quasimeromorphic mappings exist was originally posed by Martio and Srebro in [MS1]; subsequently in [MS2] they proved the existence of fore-mentioned mappings in the case of co-finite groups i.e. groups such that $Vol_{hyp}(\mathbb{H}^n/G) < \infty$ (the important case of geometrically finite groups being thus included). Also, it was later proved by Tukia ([Tu]) that the existence of non-constant quasimeromorphic mappings (or qm-maps, in short) is assured in the case when G acts torsionless upon \mathbb{H}^n . Moreover, since for torsionless Kleinian groups G , \mathbb{H}^n/G is a (analytic) manifold, the next natural question to ask is whether there exist qm-maps $f : M^n \rightarrow \widehat{\mathbb{R}^n}$; where M^n is an orientable n -manifold. The affirmative answer to this question is due to K. Peltonen (see [Pe]); to be more precise she proved the existence of qm-maps in the case when M^n is a connected, orientable \mathcal{C}^∞ -Riemannian manifold.

In contrast with the above results it was proved by Srebro ([Sr]) that, for any $n \geq 3$, there exists a Kleinian group $G \curvearrowright \mathbb{H}^n$ s.t. there exists no non-constant, G -automorphic function $f : \mathbb{H}^n \rightarrow \mathbb{R}^n$. More precisely, if G (as above) contains elliptics of unbounded orders (with non-degenerate fixed set), then G admits no non-constant G -automorphic qm-mappings.

Since all the existence results were obtained in constructive manner by using the classical "Alexander trick" (see [Al]), it is only natural that we try to attack the problem using the same method. For this reason we present here in succinct manner Alexander's method: One starts by constructing a suitable triangulation (Euclidean or hyperbolic) of \mathbb{H}^n . Since \mathbb{H}^n is orientable, an orientation consistent with the given triangulation (i.e. such that two given n -simplices having a $(n-1)$ -dimensional face in common will have opposite orientations) can be chosen. Then one quasiconformally maps the simplices of the triangulation into $\widehat{\mathbb{R}^n}$ in a chess-table manner: the positively oriented ones onto the interior of the standard simplex in \mathbb{R}^n and the negatively oriented ones onto its exterior. To ensure the existence of such a chessboard triangulation, a further barycentric type of subdivision may be required, rendering a triangulation whose simplices satisfy the condition that every $(n-2)$ -face is incident to an even number of n -simplices. If the dilatations of the qc-maps constructed above are uniformly bounded, then the resulting map will be quasimeromorphic.

The dilatations of each of the qc-maps above is dictated by the proportions of the respective simplex (see [Tu], [MS2]), and since the dilatation is to be uniformly bounded, we are naturally directing our efforts in the construction of a **fat** triangulation, where:

Definition 1.7. Let $\tau \subset \mathbb{R}^n$; $0 \leq k \leq n$ be a k -dimensional simplex. The *fatness* φ of τ is defined as being:

$$(1.3) \quad \varphi = \varphi(\tau) = \inf_{\substack{\sigma < \tau \\ \dim \sigma = l}} \frac{Vol(\sigma)}{diam^l \sigma}$$

The infimum is taken over all the faces of τ , $\sigma < \tau$, and $Vol_{eucl}(\sigma)$ and $diam \sigma$ stand for the Euclidian l -volume and the diameter of σ respectively. (If $\dim \sigma = 0$, then $Vol_{eucl}(\sigma) = 1$, by convention.)

A simplex τ is φ_0 -*fat*, for some $\varphi_0 > 0$, if $\varphi(\tau) \geq \varphi_0$. A triangulation (of a submanifold of \mathbb{R}^n) $\mathcal{T} = \{\sigma_i\}_{i \in \mathbf{I}}$ is φ_0 -*fat* if all its simplices are φ_0 -fat. A triangulation $\mathcal{T} = \{\sigma_i\}_{i \in \mathbf{I}}$ is *fat* if there exists $\varphi_0 > 0$ such that all its simplices are φ_0 -fat.

Remark 1.8. There exists a constant $c(k)$ that depends solely upon the dimension k of τ s.t.

$$(1.4) \quad \frac{1}{c(k)} \cdot \varphi(\tau) \leq \min_{\substack{\sigma < \tau \\ \dim \sigma = l}} \angle(\tau, \sigma) \leq c(k) \cdot \varphi(\tau),$$

and

$$(1.5) \quad \varphi(\tau) \leq \frac{Vol(\sigma)}{diam^l \sigma} \leq c(k) \cdot \varphi(\tau);$$

where $\angle(\tau, \sigma)$ denotes the (*internal*) *dihedral angle* of $\sigma < \tau$. (For a formal definition, see [CMS], pp. 411-412, [Som].)

Remark 1.9. The definition above is the one introduced in [CMS]. For equivalent definitions of fatness, see [Ca1], [Ca2], [Pe], [Tu].

The idea of the proof of Theorem 1.6. is, in a nutshell, as follows: first build two fat triangulations: \mathcal{T}_1 of a certain closed neighbourhood N_e^* of the singular set of \mathbb{H}^n/G ; and \mathcal{T}_2 of $(\mathbb{H}^n/G) \setminus N_e^*$; and then "mash" the two triangulations into a new triangulation \mathcal{T} , while retaining their fatness. The lift $\tilde{\mathcal{T}}$ of \mathcal{T} to \mathbb{H}^n represents the required G -invariant fat triangulation.

Since $(\mathbb{H}^n/G) \setminus N_e^*$ is an analytical manifold, the existence of \mathcal{T}_2 is assured by Peltonen's result. In [S2] we showed how to build \mathcal{T}_1 using a generalization of a theorem of Munkres ([Mun], 10.6) on extending the triangulation of the boundary of a manifold (with boundary) to the whole manifold. Munkres' technique also provided us with the basic method of mashing the triangulations \mathcal{T}_1 and \mathcal{T}_2 . In this paper we present a more direct, geometric method of triangulating N_e^* and mashing the two triangulations. We already employed this simpler method in [S1], where we proved Theorem 1.6. in the case $n = 3$. The original technique used in [S1] for fattening the intersection of \mathcal{T}_1 and \mathcal{T}_2 is, however, restricted to dimension 3. Therefore here we make appeal to the method employed in [S2], which is essentially the one developed in [CMS].

This paper is organized as follows: in Section 2 we present some background on elliptic transformations and we show how to choose and triangulate the closed neighbourhood N_e^* of the singular set of \mathbb{H}^n/G , and how to select the "intermediate zone" where the two different triangulations overlap. Section 3 is dedicated to the main task of fattening the common triangulation. In Section 4 we show how to apply the main result in the construction of a quasimeromorphic mappings from \mathbb{H}^n to $\widehat{\mathbb{R}^n}$.

2. CONSTRUCTING AND INTERSECTING TRIANGULATIONS

2.1. Elliptic Transformations. Let us first recall the basic definitions and notations: A transformation $f \in Isom(H^n)$, $f \neq Id$ is called *elliptic* if $(\exists) m \geq 2$ s.t. $f^m = Id$, and the smallest m satisfying this condition is called the *order* of f . In the 3-dimensional case the *fixed point set* of f , i.e. $Fix(f) = \{x \in H^3 | f(x) = x\}$, is a hyperbolic line and will be denoted by $A(f)$ – the *axis* of f . In dimension $n \geq 4$ the fixed set of an elliptic transformation is a k -dimensional hyperbolic plane, $0 \leq k \leq n - 2$. The situation is complicated further by the fact that different elliptics may well have fixed loci of different dimensions (inclusive the degenerate case $n = 0$). If A is an axes of an elliptic transformation of order m acting upon \mathbb{H}^3 , then A is called an m – *axis*. By extension we shall call the k -dimensional fixed point set of an elliptic transformation acting upon H^n , $n \geq 4$, an (m, k) -axis, $0 \leq k \leq n - 2$, or just an axis.

Remark 2.1. Since our main interest lies in Kleinian groups acting upon \mathbb{H}^3 , whose elliptic elements have orders bounded from above, it should be mentioned that, while for any finitely generated Kleinian group acting on H^3 the number of conjugacy classes of elliptic elements is finite (see [FM]), this is not true for Kleinian groups acting upon H^n , $n \geq 4$; (for counterexamples, see [FM], [Po] and [H]).

If the discrete group G is acting upon \mathbb{H}^n , then by the discreteness of G , there exists no accumulation point of the elliptic axes in \mathbb{H}^n . Moreover, if G contains no elliptics with intersecting axes, then the distances between the axes are, in general, bounded from below. In the classical case $n = 3$ this bounds are obtained by applying Jørgensen's inequality and its corollaries. While no actual computations in higher dimension of the said bounds are known to us, their feasibility follows from the existence of various generalizations of Jørgensen's inequality (see [FH], [M], [Wa]).¹ We defer such computations for further study. The methods above do not apply for pairs of order two elliptics (see [AH], [S1]). In this case a modification of the basic construction will be required (see Section 2.3. below). In the presence of *nodes* i.e. intersections of elliptic axes, the situation is more complicated. Indeed, even in dimension 3 the actual computation of the distances between node points has been achieved only relatively recently (see [GM1], [GM2], [GMMR])².

2.2. Geometric Neighbourhoods. To produce the desired closed neighbourhood N_e^* of the singular locus of \mathbb{H}^n / G and its triangulation \mathcal{T}_1 , we start by constructing a standard neighbourhood $N_f = N(A(f))$ of the axes of each elliptic element of G such that $N_f \simeq A(f) \times I^{n-k}$, where $A(f) = \mathbb{S}^k$ and where I^{n-k} denotes the unit $(n - k)$ -dimensional interval. The construction of N_f proceeds as follows: By [Cox], Theorem 11.23. the fundamental region for the action of the stabilizer group of the axes of f , $G_f = G_{A(f)} = \{g \in G | g(x) = x\}$ is a simplex or a product a simplices. Let \mathcal{S}_f be this fundamental region. Then we can define the generalized prism³ \mathcal{S}_f^\perp , defined by translating \mathcal{S}_f in a direction perpendicular to \mathcal{S}_f , where the translation length is $dist_{hyp}(\mathcal{S}, A(f))$. It naturally decomposes into simplices (see [Som], p. 115, [Mun], Lemma 9.4). We have thus constructed an f -invariant triangulation of a prismatic neighbourhood N_f of $A(f)$. We can reduce the mesh of this triangulation

¹ For bounds in dimension 3 see [BM], [GM1], [GM2].

² See also [DM], [Med] and [S3] for a different approach.

³ or *simplotope* cf. [Som], pp. 113-115.

as much as required, while controlling its fatness by dividing it into a finite number of radial strata of equal width $\varrho = \delta/\kappa_0$, and further partition it into "slabs" of equal height h , where $\delta = \min\{\text{dist}_{hyp}(A(f), A(g)) \mid g \text{ elliptic, } g \neq f\}$. Henceforth we shall call the neighborhood thus produced, together with its fat triangulation, a *geometric neighbourhood*.

2.3. Mashing Triangulations. Since G is a discrete group, G is countable so we can write $G = \{g_j\}_{j \geq 1}$ and let $\{f_i\}_{i \geq 1} \subset G$ denote the set of elliptic elements of G . The steps in building the desired fat triangulation are as follows:

- (1) Consider the geometric neighbourhoods

$$N_i = N_{i/4} = \{x \in \mathbb{H}^3 \mid \text{dist}_{hyp}(A_i, x) < \delta/4\}$$

and

$$N'_i = N_{i,3/16} = \{x \in \mathbb{H}^n \mid \text{dist}_{hyp}(A_i, x) < 3\delta/16\},$$

with their natural fat triangulations, where $A_i = A(f_i)$ and where the choice of " $\delta/4$ " instead of " δ " in the definition of the geometric neighbourhoods N_i is dictated by the following Lemma:

Lemma 2.2. ([Rat]) *Let X be a metric space, and let $\Gamma < \text{Isom}(X)$ be a discontinuous group.*

Then, for any $x \in X$ and any $r \in (0, \delta/4)$:

$$\pi : B(x, r)/\Gamma_x \simeq B(\pi(x), \delta/4);$$

where: Γ_x is the stabilizer of x , π denotes the natural projection, $\delta = d(x, \Gamma(x) \setminus \{x\})$, and where the metric on X/Γ is given by

$$d_\Gamma([\pi(x)], [\pi(y)]) = d(\Gamma(x), \Gamma(y)); \forall x, y \in X/\Gamma.$$

We also put: $N_e = \bigcup_{i \in \mathbb{N}} N_i$.

- (2) Denote by \mathcal{T}_e the geometric triangulation of N_e described in Section 2.2. above.
- (3) Consider the following quotients: $N_e^* = (\overline{N_e} \cap \mathbb{H}^n)/G$ and $M_c = (\mathbb{H}^n/G) \setminus N_e^* = (\mathbb{H}^n \setminus \overline{N_e})/G$.
- (4) Denote by \mathcal{T}_p the fat triangulation of M_c assured by Peltonen's Theorem.
- (5) Consider also the tubes $T_i = N_{i,1/4} \setminus N_{i,3/16}$, and denote $T = \bigcup_{i \in \mathbb{N}} T_i$.
- (6) T is endowed with two triangulations: a natural triangulation \mathcal{T}_1 induced by the geometric triangulations of the tubes T_i , and \mathcal{T}_2 that is inherited from \mathcal{T}_p .
- (7) Mashing \mathcal{T}_1 and \mathcal{T}_2 and ensuring (by a further eventual barycentric type subdivision) that each $(n-2)$ -face of every n -simplex is incident to an even number of n -simplices, produces a triangulation \mathcal{T}_0 .
- (8) Denote by \mathcal{T} the fat triangulation obtained by fattening \mathcal{T}_0 .
- (9) Let $\tilde{\mathcal{T}}$ be the lift of \mathcal{T} to \mathbb{H}^n . Then $\tilde{\mathcal{T}}^\# = \tilde{\mathcal{T}} \cup \mathcal{T}_e$ represents the desired G -invariant fat triangulation of \mathbb{H}^n .

Remark 2.3. The construction above applies only in the case of non-intersecting elliptic axes. In the case when there exist intersecting elliptic axes, the following modification of our construction is required: instead of δ one has to consider $\delta^* = \min(\delta, \delta_0)$, where δ_0 represents the minimal distance between nodes.

Remark 2.4. In the case when all the elliptic transformations are half-turns, since, as we have seen, no minimal distance between the axes can be computed in this case. However, by the discreteness of G it follows that there is no accumulation point of the axes in \mathbb{H}^3 . Let $D = \{d_{ij} \mid d_{ij} = \text{dist}_{hyp}(A_i, A_j)\}$ denote the set of mutual distances between the axes of the elliptic elements of G . Then, since G is countable, so will be D , thus $D = \{d_k\}_{k \geq 1}$. Then the set of neighbourhoods $N_e^h = \bigcup_{k \in \mathbb{N}} N_k^h = \bigcup_{k \in \mathbb{N}} \{x \in \mathbb{H}^n \mid \text{dist}_{hyp}(A_k, x) < \delta/4k\}$ will constitute a proper geometric neighbourhood of $A_G = \bigcup_{i \in \mathbb{N}} A_i$. The fatness of the simplices of the geometric triangulation of A_G can be controlled, as before, by a proper choice of h and ϱ .

3. FATTENING TRIANGULATIONS

First let us establish some definitions and notations:

Let K denote a simplicial complex, let $K' < K$ denote a subcomplex of K and let $\sigma \in K$ denote the simplices of K .

Definition 3.1. Let $\sigma_i \in K$, $\dim \sigma_i = k_i$, $i = 1, 2$; s.t. $\text{diam } \sigma_1 \leq \text{diam } \sigma_2$. We say that σ_1, σ_2 are δ -transverse iff

(i) $\dim(\sigma_1 \cap \sigma_2) = \max(0, k_1 + k_2 - n)$;

(ii) $0 < \delta < \angle(\sigma_1, \sigma_2)$;

and if $\sigma_3 \subset \sigma_1$, $\sigma_4 \subset \sigma_2$, s.t. $\dim \sigma_3 + \dim \sigma_4 < n = \dim K$, then

(iii) $\text{dist}(\sigma_3, \sigma_4) > \delta \cdot \eta_1$.

In this case we write: $\sigma_1 \pitchfork_\delta \sigma_2$.

We begin by triangulating and fattening the intersection of two individual simplices belonging to the two given triangulations, respectively. Given two closed simplices $\bar{\sigma}_1, \bar{\sigma}_2$, their intersection (if not empty) is a closed, convex polyhedral cell: $\bar{\gamma} = \bar{\sigma}_1 \cap \bar{\sigma}_2$. One canonically triangulates $\bar{\gamma}$ by using the *barycentric subdivision* $\bar{\gamma}^*$ of $\bar{\gamma}$, defined inductively upon the dimension of the cells of $\partial \gamma$ in the following manner: for each cell $\beta \subset \partial \gamma$, choose an interior point $p_\beta \in \text{int } \beta$ and construct the join $J(p_\beta, \beta_i)$, $\forall \beta_i \subset \partial \beta$.⁴

We first show that if the simplices are fat and if they intersect δ -transversally, then one can choose the points s.t. the barycentric subdivision $\bar{\gamma}^*$ will be composed of fat simplices. More precisely, we prove the following Proposition:

Proposition 3.2. Let $\sigma_1, \sigma_2 \subset \mathbb{R}^m$, where $m = \max(\dim \sigma_1, \dim \sigma_2)$, s.t. $d_1 = \text{diam } \sigma_1 \leq d_2 = \text{diam } \sigma_2$, and s.t. σ_1, σ_2 have common fatness φ_0 . If $\sigma_1 \pitchfork_\delta \sigma_2$, then there exists $c = c(m, \varphi_0, \delta)$

- (1) If $\sigma_3 \subset \bar{\sigma}_1$, $\sigma_4 \subset \bar{\sigma}_2$ and if $\sigma_1 \cap \sigma_2 \neq \emptyset$, then $\sigma_1 \cap \sigma_2 = \gamma_0$ is an $(k_3 + k_4 - m)$ -cell, where $k_3 = \dim \sigma_3$, $k_4 = \dim \sigma_4$ and:

$$(3.1) \quad \text{Vol}_{\text{eucl}}(\gamma_0) \geq c \cdot d_1^{k_3 + k_4 - m}.$$

- (2) $\forall \gamma_0$ as above, $\exists p \in \gamma_0$, s.t.

$$(3.2) \quad \text{dist}(p, \partial \gamma_0) > c \cdot d_1.$$

⁴ If $\dim \beta = 0$ or $\dim \beta = 1$, then β is already a simplex.

- (3) If the points employed in the construction of $\bar{\gamma}^*$ satisfy the condition (3.2) above, then each l -dimensional simplex $\tau \in \gamma^*$ satisfies the following inequalities:

$$(3.3) \quad \varphi_l \geq Vol_{eucl}(\tau)/d_1^l \geq c \cdot d_1.$$

Proof. First, consider the following remarks:

Remark 3.3. The following sets are compact:

$$S_1 = \{\sigma_1 \mid diam \sigma_1 = 1, \varphi(\sigma_1) \geq \varphi_0\}, S_2 = \{\sigma_2 \mid diam \sigma_2 = 2(1 + \delta), \varphi(\sigma_2) \geq \varphi_0\},$$

$$S(\phi_0, \delta) \subset S_1 \cap S_2, S(\phi_0, \delta) = \{(\sigma_1, \sigma_2) \mid \exists v_0, \text{ s.t. } v_0 \in \sigma_1, \forall \sigma_1 \in S_1 \cap S_2\}.$$

Remark 3.4. There exists a constant $c(\varphi)$ s.t. $\mathcal{S} = \mathcal{S}'$, where

$$\mathcal{S} = \{\sigma_1 \cap \sigma_2 \mid diam \sigma_2 \leq d_2\}, \mathcal{S}' = \{\sigma_1 \cap \sigma_2 \mid diam c(\varphi)(1 + \delta)d_1\},$$

i.e. the sets of all possible intersections remains unchanged under controlled dilations of one of the families of simplices.

Now, from the fact that $\sigma_1 \pitchfork_\delta \sigma_2$ it follows that $\sigma_3 \cap \sigma_4 \neq \emptyset \Leftrightarrow \bar{\sigma}_3 \cap \bar{\sigma}_4 \neq \emptyset$ (see [CMS], p. 436). Therefore, the function $Vol_{eucl}(\gamma_0)$ attains a positive minimum, as a positive, continuous function defined on the compact set $\bar{\sigma}_3 \cap \bar{\sigma}_4$, thus proving the first assertion of the proposition.

Let be γ be a q -dimensional cell, and let β be a face of $\partial\gamma$. Then:

$$(3.4) \quad Vol_{eucl}(\beta) \leq d_1^p.$$

Choose $p \in \gamma$, such that $\rho = dist(p, \partial\gamma) = \max\{dist(r, \partial\gamma) \mid r \in \gamma\}$. Then, if $\beta = \beta^j$ denotes a j -dimensional face of $\partial\gamma$, we have that:

$$(3.5) \quad \gamma \subset \bigcup_{\beta^j \subset \partial\gamma} N_\rho(\beta^j);$$

where: $N_\rho(\beta^j) = \{r \mid dist(r, \beta^j) \subseteq \rho\}$. But:

$$(3.6) \quad Vol_{eucl}(\beta^j \cap \gamma) \leq c \cdot \rho^{q-j} \cdot Vol_{eucl}(\beta^j),$$

for some $c' = c'(q)$.

Moreover, the number of faces $\sigma_3 \cap \sigma_4$ of γ is at most $2^{dim \sigma_1 + dim \sigma_2 + 2}$, where σ_1, σ_2 are as in Remark 3.4. and $dim \sigma_3 \leq dim \sigma_1$, $dim \sigma_4 \leq dim \sigma_2$.

Thus (3.4) in conjunction with (3.6) imply that there exists $c_1 = c_1(m, \varphi_0, \delta)$, such that:

$$(3.7) \quad c_1 d_1^q \leq \sum_{j=0}^{q-1} \rho^{q-j} d_1^j.$$

and (3.2) follows from this last inequality.

The last inequality follows from (3.2) and (3.3) by induction. \square

Next we show that given two fat Euclidian triangulations that intersect δ -transversally, then one can infinitesimally move any given point of one of the triangulations s.t. the resulting intersection will be δ' -transversal, where δ' depends only on δ , the

common fatness of the given triangulations, and on the displacement length. More precisely one can show that the following results holds:

Proposition 3.5. *Let $K_1, K_2 \subset \mathbb{R}^n$ be n -dimensional simplicial complexes, of common fatness φ_0 and $d_1 = \text{diam } \sigma_1 \leq d_2 = \text{diam } \sigma_2$. Let $v_0 \in K_1$ be a 0-dimensional simplex of K_1 . Consider the complex K_1^* obtained by replacing v_0 by $v_0^* \in \mathbb{R}^n$ and keeping fixed the rest of the 0-dimensional vertices of K_1 fixed. Consider also $L_2 < K_2$, $L_2 = \{\sigma \in K_2 \mid \sigma \cap B(v_0, 2d_1) \neq \emptyset\}$. Then, if there exists k s.t. all the k -simplices $\tau \subset \partial St(v_0)$ are δ -transversal to L_2 , there exist $\varphi_0, \delta, \varepsilon > 0$, $\delta^* = \delta^*(\varphi_0, \delta, \varepsilon)$ and there exists v_0^* s.t. $\text{dist}(v_0, v_0^*) < \varepsilon \cdot d_1$ s.t.*

$$(3.8) \quad \tau^* \pitchfork_{\delta^*} L_2; \forall \tau \subset St(v_0^*) \setminus \partial St(v_0^*), \dim \tau^* = k + 1.$$

Proof. Let $N(r) = |\{\sigma \in K_1 \mid \sigma \subset B_r(v_0)\}|$. Then there exists a constant c_n s.t. $N(r) \leq \frac{c_n}{\varphi_0} (\frac{\varepsilon}{d_1})^n$. It follows that the set $St(v_0)$ is compact, since there are at most $\frac{c_n}{\varphi_0}$ possible edge lengths, which can take values in the interval $[d_1 \varphi_0, d_1]$.⁵ Therefore if a D^* satisfying (3.8) exist, it depends only on φ_0 , δ and ε (and not on K_1, K_2). Let $\sigma_1, \dots, \sigma_{l_1}$ and $\tau_1, \dots, \tau_{l_2}$ be orderings of the simplices of L_2 and of the k -simplices of $\partial St(v_0)$, respectively. Then, by [CMS], Lemma 7.4, there exists $\varepsilon_{1,1}$ and exists $v_{1,1}$, $d(v_{1,1}, v_0) = \varepsilon_{1,1}$, such that the hyperplane $\Pi(v_{1,1}, \tau_1)$ determined by v_1 and by τ_1 is transversal to σ_1 . By replacing τ_1 by τ_2 and v_0 by $v_{1,1}$ we obtain $v_{1,2}$ and $\varepsilon_{1,2}$ s.t. $\Pi(v_{1,2}, \tau_1) \pitchfork \sigma_2$. (See Fig. 1.)

Moreover, by choosing $\varepsilon_{1,2}$ sufficiently small, one can ensure that $\Pi(v_{1,2}, \tau_1) \pitchfork \sigma_1$, also. Repeating the process for $\tau_3, \dots, \tau_{l_2}$, one determines a point v_{1,l_2} such that $\Pi(v_{1,l_2}, \tau_j) \pitchfork L_2$ $j = 3, \dots, l_2$. In the same manner and by choosing at each stage an $\varepsilon_{i,j}$ small enough, one finds points $v_{i,j}$ s.t. $\Pi(v_{i,j}, \tau_i) \pitchfork L_2$, $i = 1, \dots, l_1$, $j = 1, \dots, l_2$. Then $v_0^* = v_{l_1, l_2}$ satisfies: $\Pi(v_0^*, \tau_j) \pitchfork L_2$, $j = 1, \dots, l_2$. \square

We are now prepared to prove the main result of this section namely:

Theorem 3.6. *Let M^n be an orientable n -manifold and let $\mathcal{T}_1, \mathcal{T}_2$ be two fat triangulations of open sets $U_1, U_2 \subset M^n$, $U_1 \cap U_2 \neq \emptyset$, having common fatness $\geq \varphi_0$, and such that $\mathcal{T}_1 \cap \mathcal{T}_2 \neq \emptyset$. Then there exist fat triangulations $\widetilde{\mathcal{T}}'_1, \mathcal{T}'_2$ and there exist open sets $U \subset U_1 \cap U_2 \subset V$, such that*

- (1) $(\mathcal{T}'_1 \cap \mathcal{T}'_2) \cap (U_i \setminus V) = \mathcal{T}_i$, $i = 1, 2$;
- (2) $(\mathcal{T}'_1 \cap \mathcal{T}'_2) \cap U = \mathcal{T}$;

where

- (3) \mathcal{T} is a fat triangulation of U .

Proof. Let K_1, K_2 denote the underlying complexes of $\mathcal{T}_1, \mathcal{T}_2$, respectively. By considerations similar to those of Proposition 3.5. it follows that given $\varphi_0 > 0$, there exists $d(\varphi_0) > 0$ such that given a k -dimensional simplex $\sigma \subset \mathbb{R}^n$, $\text{diam}(\sigma) = d_1$ has fatness φ_0 , then translating each vertex of σ by a distance $d(\varphi_0) \cdot d_1$ renders a simplex of fatness $\geq \varphi_0/2$. Also, it follows that given $\varphi_0, \delta > 0$, exists $\delta(\varphi_0, \delta)$ satisfying the following condition: if every vertex $u \in \sigma \subset K$ is replaced by a vertex u' s.t. $\text{dist}(u, u') \leq \delta(\varphi_0, \delta) \cdot d_1$, then the resulting simplex σ' is $\pitchfork_{\delta/2}$ -transversal to K ; for any n -dimensional simplicial complex K of fatness φ_0 and such that

⁵ i.e. the number of possible combinatorial structures on $St(v_0)$ depends only on φ_0 .




FIGURE 1.

$\text{diam } \sigma = d_2 \geq d_1$.

Let $v_0 \in U_1 \cap U_2$. Define the following subcomplexes of K_1 , K_2 , respectively:

$$L_2 = \{\bar{\sigma} \subset K_2 \mid \bar{\sigma} \subset B_\varepsilon(v_0), d_1 \leq \text{dist}(\bar{\sigma}, \partial B_\varepsilon(v_0)) \leq d_2\}$$

$$M_2 = \{\sigma \subset K_2 \mid \bar{\sigma} \subset \bar{\tau} \subset B_\varepsilon(v_0), \dim \tau = n, \bar{\sigma} \cap L_2 \neq \emptyset\}$$

$$L_1 = \{\bar{\sigma} \subset K_1 \mid \text{dist}(\bar{\sigma}, L_2) \leq d_2\}$$

$$M_1 = \{\sigma \subset K_1 \mid \bar{\sigma} \subset \bar{\tau} \subset B_\varepsilon(v_0), \dim \tau = n, \tau \cap L_1 \neq \emptyset\}$$

(See Fig. 2.)

Consider an ordering v_1, \dots, v_p of the vertices of L_1 . It follows from Proposition 3.5. that, if all the vertices of L_1 are moved by at most t_0 , where

$$(3.9) \quad t_0 = \frac{d_1}{n} \min \left\{ \frac{1}{2}, d(\varphi_0) \right\},$$

then there exists

$$(3.10) \quad \delta_0^* = \delta_0^*(\varphi_0, 1, \frac{t_0}{d_1}),$$

such that

$$(3.11) \quad \mathcal{S}^0(L_{1,0}) \pitchfork_{\delta_0^*} K_2,$$

where $\mathcal{S}^0(L_{1,0})$ denotes the 0-skeleton of $L_{1,0}$.




FIGURE 2.

Now define inductively

$$(3.12) \quad t_i = \frac{d_1}{n} \min \left\{ \frac{1}{2}, d(\varphi_0), \delta\left(\frac{\varphi_0}{2}, \frac{\delta_0^*}{2}\right), \dots, \delta\left(\frac{\varphi_0}{2}, \frac{\delta_{i-1}^*}{2}\right) \right\},$$

where

$$(3.13) \quad \delta_i^* = \delta_i^*\left(\varphi_0, \delta_{i-1}^*, \frac{t_i}{d_1}\right); \quad i = 1, \dots, n-1.$$

Then $t_0 \geq t_1 \geq \dots \geq t_{n-1}$.

Moving each and every vertex of L_1 by a distance $\leq t_i$, $i = 1, \dots, n$, renders complexes $L_{1,1}, \dots, L_{1,n-1}$ s.t.

- (1) $L_{1,i} \cap (B_\varepsilon(v_0) \setminus M_1) \equiv L_1$,
- (2) $L_{1,i}$ are φ_0 -fat; $i = 0, \dots, n-1$.

By inductively applying Proposition 3.5. it follows that

$$(3.14) \quad \mathcal{S}^i(L_{1,i}) \pitchfork_{\delta_i^*} K_2,$$

Where $\mathcal{S}^i(L_{1,i})$ denotes the i -skeleton of $L_{1,i}$.
Moreover,

$$(3.15) \quad \mathcal{S}^i(L_{1,j}) \pitchfork_{\delta_i^*/2} K_2, \forall j > i.$$

It follows that

$$(3.16) \quad L_{1,n-1} \pitchfork_{\delta^*} K_2,$$

where

$$(3.17) \quad \delta^\star = \frac{1}{2} \min\{\delta_0^*, \dots, \delta_i^*\}.$$

By Proposition 3.2. the barycentric subdivision of $L_{1,n-1} \cap L_2$ is fat. We extend it to a fat subdivision of M_2 in the following manner: given a simplex $\sigma \subset M_2 \setminus L_2$, subdivide σ by constructing all the simplices with vertices v_i , where v_i is either the vertex of a simplex $\sigma \subset M_2 \setminus L_2$, $\bar{\sigma}_i \cap L_2 \neq \emptyset$, or it is a vertex of a closed simplex $\bar{\sigma}$ of the barycentric subdivision of $L_{1,n-1} \cap L_2$, such that $\bar{\sigma} \subset \partial L_2 \cap M_2$, $i = 1, \dots, k_0$. The triangulation \tilde{K}_2 thus obtained is a fat extension of $\overline{K_2 \setminus M_2}$.

In an analogous manner one constructs a similar fat extension \tilde{K}_1 of $\overline{K_1 \setminus M_2}$. \square

Now let $M^n = M_c$ and let $\mathcal{T}_1, \mathcal{T}_2$ be the fat chessboard triangulations constructed in 2.3.(6)–2.3.(7). Then the local fat triangulation obtained in Theorem 3.6. above extends globally to the fat triangulation of $\mathcal{T}_1 \cap \mathcal{T}_2$, by applying Lemma 10.2 and Theorem 10.4 of [Mun]. This provides us with the sought fat chessboard triangulation \mathcal{T} of M_c . Thus $\tilde{\mathcal{T}}^\sharp = \tilde{\mathcal{T}} \cup \mathcal{T}_e$ represents the required G -invariant fat chessboard triangulation of \mathbb{H}^n .

4. THE EXISTENCE OF QUASIMEROMORPHIC MAPPINGS

The technical ingredient in Alexander's trick is the following Lemma:

Lemma 4.1. ([MS1], [Pe]) *Let M^n be an orientable n -manifold, let \mathcal{T} be a chessboard fat triangulation of M^n and let $\tau, \sigma \in \mathcal{T}$, $\tau = (p_1, \dots, p_n)$, $\sigma = (q_1, \dots, q_n)$; and denote $|\tau| = \tau \cup \text{int } \tau$.*

Then there exists a orientation-preserving homeomorphism $h = h_\tau : |\tau| \rightarrow \widehat{\mathbb{R}^n}$ s.t.

- (1) $h(|\tau|) = |\sigma|$, if $\det(p_1, \dots, p_n) > 0$
and
 $h(|\tau|) = \widehat{\mathbb{R}^n} \setminus |\sigma|$, if $\det(p_1, \dots, p_n) < 0$.
- (2) $h(p_i) = q_i$, $i = 1, \dots, n$.
- (3) $h|_{\partial|\sigma|}$ is a PL homeomorphism.
- (4) $h|_{\text{int}|\sigma|}$ is quasiconformal.

Proof Let $\tau_0 = (p_{0,1}, \dots, p_{0,n})$ denote the equilateral n -simplex inscribed in the unit sphere \mathbb{S}^{n-1} . The radial linear stretching $\varphi : \tau \rightarrow \overline{B^n}$ is onto and bi-lipschitz (see [MS2]). Moreover, by a result of Gehring and Väisälä, φ is also quasiconformal (see [V]). We can extend φ to $\widehat{\mathbb{R}^n}$ by defining $\varphi(\infty) = \infty$. Let J denote the reflection in the the unit sphere \mathbb{S}^{n-1} and let $h_0 : |\sigma| \rightarrow |\tau|$ denote the orientation-reversing PL mapping defined by: $h(p_i) = q_i$, $i = 1, \dots, n$. Then $h = \varphi^{-1} \circ J \circ \varphi \circ h_0$ is the required homeomorphism. \square

The Existence Theorem of quasimeromorphic mappings now follows immediately:

Proof of Theorem 1.6. Since the orders of the elliptic transformations are uniformly bounded, so will be the fatness of the simplices of \mathcal{T}_1 – the geometric triangulation of T . Let $\tilde{\mathcal{T}}^\sharp$ be the G -invariant fat chessboard triangulation of \mathbb{H}^n constructed above. Let $f : \mathbb{H}^n \rightarrow \widehat{\mathbb{R}^n}$ be defined by: $f|_{|\sigma|} = h_\sigma$, where h is

the homeomorphism constructed in the Lemma above. Then f is a local homeomorphism on the $(n-1)$ -skeleton of \tilde{T}^\sharp too, while its branching set B_f is the $(n-2)$ -skeleton of \tilde{T}^\sharp . By its construction f is quasiregular. Moreover, given the uniform fatness of the simplices of triangulation \hat{T}^\sharp , the dilatation of f depends only on the dimension n . □

REFERENCES

- [Abi] Abikoff, W.: *Kleinian Groups*, Lecture Notes, given at "The Technion – Israel Institute of Technology", Haifa, Israel, 1996-1997.
- [AH] Abikoff, W. and Hass, A.H.: *Nondiscrete groups of hyperbolic motions*, Bull. London Math. Soc. 22, 1990, no.3, pp. 233-238.
- [Al] Alexander, J.W.: *Note on Riemann spaces*, Bull. Amer. Math. Soc. 26, 1920, pp. 370-372.
- [AP1] Apanasov, B.N.: *Klein Groups in Space*, Sib. Math. J. 16, 1975, pp. 679-684.
- [AP2] Apanasov, B.N.: *Discrete groups in space and uniformization problems*, Kluwer Academic Publishers, Dordrecht, 1991.
- [BM] Bowditch, B.H. and Mess, G.: *A 4-Dimensional Kleinian Group*, Transaction of the Amer. Math. Soc., Vol. 344, No.1, 1994, pp. 390-405.
- [BrM] Brooks, R. and Matelski, J.P.: *Collars in Kleinian groups*, Duke Math.J., Vol. 49, No. 1, 1982, pp. 163-182.
- [Ca1] Cairns, S.S.: *On the triangulation of regular loci*, Ann. of Math. 35, 1934, pp. 579-587.
- [Ca2] Cairns, S.S.: *Polyhedral approximation to regular loci*, Ann. of Math. 37, 1936, pp. 409-419.
- [CMS] Cheeger, J., Müller, W., and Schrader, R.: *On the Curvature of Piecewise Flat Spaces*, Comm. Math. Phys., 92, 1984, 405-454.
- [Cox] Coxeter, H. S. M.: *Regular Polytopes*, Second Edition, MacMillan, London, 1963.
- [DM] Derevin, D.A. and Mednikh, A.D.: *Geometric proprieties of discrete groups acting with fixed points in Lobachevsky space*, Soviet Math. Dokl., Vol. 37, 1988, No.3, pp. 614-617.
- [FH] Friedland, S. and Hersonsky, S.: *Jorgensen's inequality for discrete groups in normed algebras*, Duke Mathematical Journal, Vol. 69, No. 3, 1993.
- [FM] Feighn, M. and Mess, G.: *Conjugacy classes of finite subgroups of Kleinian groups*, Amer. J. of Math., 113, 1991, pp. 179-188.
- [GM1] Gehring, F.W., Martin G.J., *Commutators, collars and the geometry of Möbius groups*, J. Anal. Math. Vol. 63, 1994, pp. 174-219.
- [GM2] Gehring, F.W., Martin, G.J., *On the Margulis constant for Kleinian groups, I*, Ann. Acad. Sci. Fenn., Vol. 21, 1996, pp. 439-462.
- [GMMR] Gehring, F.W., MacLachlan, C., Martin, G.J., and Reed, A.W.: *Aritmecity, Discreteness and Volume*, Trans. Amer. Math. Soc. 349, 1997, 3611-3643.
- [H] Hamilton, E.: *Geometrical finiteness for hyperbolic orbifolds*, Topology, Vol.37, No.3, 1998, pp. 635-657.
- [J] Jørgensen, T.: *On discrete groups of Möbius transformations*, Amer. Journ. of Math., Vol. 98, No.3, pp.739-749.
- [M] Martin, G.J.: *On discrete Möbius groups in all dimensions: A generalization of Jørgensen's inequality*, Acta Math. 163, 1989, pp. 253-289.
- [Med] Mednikh, A.D.: *Automorphism groups of the three-dimensional hyperbolic manifolds*, Soviet Math. Dokl., Vol. 32, 1985, No.3, pp. 633-636.
- [MS1] Martio, O., and Srebro, U.: *Automorphic quasimeromorphic mappings in R^n* , Acta Math. 195, 1975, pp. 221-247.
- [MS2] Martio, O., and Srebro, U.: *On the existence of automorphic quasimeromorphic mappings in R^n* , Ann. Acad. Sci. Fenn., Series I Math., Vol. 3, 1977, pp. 123-130.
- [Ms] Maskit, B.: *Kleinian Groups*, Springer Verlag, GDM 287, N.Y., 1987.
- [Mun] Munkres, J. R.: *Elementary Differential Topology*, (rev. ed.) Princeton University Press, Princeton, N.J., 1966.
- [Pe] Peltonen, K.: *On the existence of quasiregular mappings*, Ann. Acad. Sci. Fenn., Series I Math., Dissertationes, 1992.
- [Po] Potyagailo, L.: *Finitely generated Kleinian groups in 3-space and 3-manifolds of infinite homotopy type*, Trans. of Amer. Math. Soc., Vol. 344, No. 1, 1994, pp. 57-77.

- [Rat] Ratcliffe, J.G.: *Foundations of Hyperbolic Manifolds*, GTM 194, Springer Verlag, N.Y., 1994.
- [S1] Saucan, E.: *The Existence of Quasimeromorphic Mappings in Dimension 3*, submitted to "Conformal Geometry and Dynamics".
- [S2] Saucan, E.: *Note on a theorem of Munkres*, submitted to "The Asian Journal of Mathematics".
- [S3] Saucan, E.: *On the Existence of Quasimeromorphic Mappings*, Ph. Thesis, Technion, in preparation.
- [Som] Sommerville, D.M.Y.: *An introduction to the Geometry of N Dimensions*, Dover Publications, N.Y., 1958.
- [Sr] Srebro, U. : *Non-existence of Automorphic Quasimeromorphic Mappings*, Analysis and topology, World Sci. Publishing, River Edge, NJ, 1998.
- [Th] Thurston, W.: *Three-Dimensional Geometry and Topology, vol.1, (Edited by S. Levy)*, Princeton University Press, Princeton, N.J. 1997.
- [Tu] Tukia, P.: *Automorphic Quasimeromorphic Mappings for Torsionless Hyperbolic Groups*, Ann. Acad. Sci. Fenn., 10, 1985, pp. 545-560.
- [V] Väisälä, J.: *Lectures on n-dimensional quasiconformal mappings*, Lecture Notes in Mathematics 229, Springer-Verlag, Berlin - Heidelberg - New-York, 1971.
- [Wa] Waterman, P.L.: *Möbius Transformations in Several Dimensions*, Advances in Mathematics, 101, 1993, pp. 87-113.

DEPARTMENT OF MATHEMATICS, TECHNION, HAIFA AND SOFTWARE ENGINEERING DEPARTMENT,
ORT BRAUDE COLLEGE, KARMIEL, ISRAEL

E-mail address: `semil@tx.technion.ac.il`